

# Almost tetrahedral manifolds

Evgeny Fominykh

St. Petersburg State University

6-th International Conference  
GROUPS AND QUANDLES IN LOW-DIMENSIONAL TOPOLOGY  
June 11, 2025  
Tomsk

## Ideal triangulations of 3-manifolds with boundary

Let  $M$  be a compact connected 3-manifold with non-empty boundary

An **ideal tetrahedron** is a tetrahedron with its vertices removed.


An **ideal triangulation** of  $M$  is a realization of the interior of  $M$  as a gluing of some **ideal tetrahedra**, induced by a simplicial pairing of the faces.

B.G. Casler, 1965

Every  $M$  has an ideal triangulation.



$ABD - FEH$ ,  $BCD - EFG$ ,  
 $ADC - EGH$ ,  $ACB - FHG$



A triangulation of  $M$  is **minimal** if there is no triangulation of  $M$  with fewer tetrahedra.

The number of tetrahedra in a minimal triangulation of  $M$  is denoted  $c_{\Delta}(M)$  and termed the **triangulation complexity** of  $M$ .

**Problem.**

How to find the triangulation complexity of a given 3-manifold.

V. Turaev – A. Vesnin – E. F., 2016

If  $\mathcal{T}$  has exactly **two** edges, and  $\mathcal{T}$  does not admit a 3-2 Pachner move, then  $\mathcal{T}$  is minimal.

A. Ryabkov – E. Shumakova – E. F.,

Minimality of ideal triangulations with exactly **three** edges.

D. Nigomedyanov – E. F., 2023


Let  $M$  be a connected compact 3-manifold with boundary. Then

$$c_{\Delta}(M) \geq \beta_1(M, \mathbb{Z}_2).$$

- Low bounds can be obtained from the information about the hyperbolic volume.


$$c_{\Delta}(M) \geq \frac{\text{vol}(M)}{v_3},$$

- $v_3 = 1.01494\dots$
- $v_3$  is the volume of the regular ideal hyperbolic tetrahedron;
- every geodesic tetrahedron in  $\mathbb{H}^3$  has volume at most  $v_3$ .



## When the volume does not help.

Every twist knot  $K_n$  (depicted on the right) can be obtained by Dehn filling the red component of the Whitehead link  $L$  (depicted on the left).



$vol(S^3 \setminus L) = v_{oct} = \text{volume of a regular ideal octahedron in } \mathbb{H}^3 = 3.6638\dots$

**Theorem (M. Gromov and W. Thurston)**

Let  $M$  be a finite volume hyperbolic manifold with cusps. Let  $N$  be a Dehn filling of some cusps of  $M$ . Then  $vol(N) < vol(M)$ .

$$vol(S^3 \setminus K_n) < v_{oct}$$

We call a cusped finite volume hyperbolic 3-manifold  $M$  **tetrahedral** if it can be decomposed into **regular ideal tetrahedra**.

### Theorem.

If the number of tetrahedra is  $k$ , then  $c_\Delta(M) = k$ .

Proof:

- Since  $M$  is obtained by gluing  $k$  ideal tetrahedra, we have  $c_\Delta(M) \leq k$ .
- $c_\Delta(M) \geq \frac{\text{vol}(M)}{v_3}$ , where  $v_3 = 1.01494\dots$  is the volume of the regular ideal tetrahedron.
- $c_\Delta(M) \geq k$ , since  $\text{vol}(M) = k \cdot v_3$ .

- [SnapPy](#) is a modern user interface to the Jeff Weeks's SnapPea kernel. SnapPy combines a link editor and 3D-graphics for Dirichlet domains and cusp neighborhoods with a powerful command-line interface based on the Python programming language. Can be used under the [Sage](#). Project by Marc Culler and Nathan Dunfield.
- [3-Manifold Recognizer](#) accepts many different presentations of 3-manifolds, calculates their different invariants and in many cases completely recognizes them. Project by Sergei Matveev, Vladimir Tarkaev and Chelyabinsk topology group.

Burton – Callahan – Hildebrand – Thistlethwaite – Weeks census

Tetrahedra	Orientable manifolds	
	Total	Tetrahedral
1	0	0
2	2	2
3	9	0
4	56	4
5	234	2
6	962	7
7	3 552	1
8	12 846	13
9	44 250	1
Total	61 911	30

[S. Garoufalidis – M. Goerner – V. Tarkaev – A. Vesnin – E.F., 2016

There exists **11,580** orientable tetrahedral manifolds up to **25** tetrahedra.

There exists **25,194** non-orientable tetrahedral manifolds up to **21** tetrahedra.

### Remark.

If  $N$  is a  $k$ -fold covering of a tetrahedral manifold  $M$ , then  $N$  is also tetrahedral and

$$c_{\Delta}(N) = k \cdot c_{\Delta}(M).$$

This gives infinite families of manifolds with known complexity.

**Example:** Let  $N_k$  be the total space of the punctured torus bundle over  $S^1$  with monodromy  $\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}^k$ .

[S. Anisov, 2005]


$$c_{\Delta}(N_k) = 2k.$$

Proof:  $N_k$  is the  $k$ -fold covering of the figure-eight knot complement  $N_1$ .


We say that a triangulation of a cusped finite volume hyperbolic 3-manifold  $M$  with  $k$  tetrahedra is **almost tetrahedral** if

$$(k - 1) \cdot v_3 < \text{vol}(M) < k \cdot v_3.$$

If such a triangulation exists, we say  $M$  is **almost tetrahedral**.



Whitehead link (L5a1) complement ( $m129$ )  
4 tetrahedra,  $\text{vol} = 3.66386237671\dots$



Borromean rings (L6a4) complement ( $t12067$ )  
8 tetrahedra,  $\text{vol} = 7.32772475342\dots$


## Burton – Callahan – Hildebrand – Thistlethwaite – Weeks census

Tetrahedra	Orientable manifolds		
	Total	Tetrahedral	Almost Tetrahedral
1	0	0	0
2	2	2	0
3	9	0	9
4	56	4	50
5	234	2	144
6	962	7	358
7	3 552	1	675
8	12 846	13	1 467
9	44 250	1	3 239
Total	61 911	30	5 942

### Theorem.

If  $T$  is an almost tetrahedral decomposition of  $M$  with  $k$  tetrahedra, then  $c_{\Delta}(M) = k$ .

## Example: complements of 2-bridge knots and links.



We can represent a two-bridge link  $K(p/q)$  by using Conway's notation as

$$p/q = [a_1, a_2, \dots, a_{n-1}, a_n] = a_1 + \frac{1}{a_2 + \dots + \frac{1}{a_{n-1} + \frac{1}{a_n}}}.$$

Here  $a_j$  denotes a number of half-twists.

[M. Ishikawa – K. Nemoto, 2016]

If  $p/q = [2, 1, 1, \dots, 1, 2]$ , then  $c_{tet}(S^3 \setminus K(p, q)) = 2n - 2$ .

Proof:

- [M. Sakuma – J. Weeks, 1995] and [M. Ishikawa – K. Nemoto, 2016]: constructed ideal triangulations of  $S^3 \setminus K(p, q)$  with  $2n - 2$  tetrahedra.
- [C. Petronio – A. Vesnin, 2009] based on [D. Futer – E. Kalfagianni – J. Purcell, 2008]:

$$\text{vol}(S^3 \setminus K(p, q)) > (2n - 2.66) \cdot v_3.$$

Let  $M$  be a finite volume hyperbolic 3-manifold. Let  $S$  be an incompressible thrice-punctured sphere properly embedded in  $M$ . Then  $S$  is isotopic to a thrice-punctured sphere  $S'$  properly embedded in  $M$  such that  $S'$  is totally geodesic in the hyperbolic structures on  $M$ .

Let  $S$  be an incompressible thrice-punctured sphere in an orientable finite volume hyperbolic 3-manifold  $M$ . Let  $M'$  be the 3-manifold obtained by cutting  $M$  open along  $S$  and then reidentifying the two copies of  $S$  by an orientation-preserving homeomorphism of  $S$ . Then  $M'$  is hyperbolic with the same volume as  $M$ .

One can cut two finite volume hyperbolic 3-manifolds  $M_1$  and  $M_2$  open along embedded incompressible thrice-punctured spheres  $S_1$  and  $S_2$  contained in  $M_1$  and  $M_2$ , respectively, and then glue copies of the thrice-punctured spheres together to yield a hyperbolic 3-manifold  $M$  with volume equal to the sum of the volumes of  $M_1$  and  $M_2$ .

## How to construct a new almost tetrahedral manifold


### Example

If  $M$  is a finite volume hyperbolic 3-manifold with an ideal triangulation  $T_M$  containing an **embedded thrice-punctured sphere**  $S_M$  which is realized by **two faces of the triangulation**, then  $S_M$  is suitable for the Adams method.


Tetrahedra	Tetrahedral	Almost Tetrahedral
4	0	2
5	2	0
6	0	9
7	0	23
8	0	16
9	0	63
10	29	?

tetrahedral mfd + almost tetrahedral mfd = new almost tetrahedral mfd

We switch from the viewpoint of ideal triangulations to the dual viewpoint of standard polyhedra.



genuine cell structure



simple polyhedron



*Nonsingular point*




*Triple line*




*True vertex*


dual polyhedron  $P$  of triangulation

- simple polyhedron;
- genuine cell structure;
- $P$  is homotopy equiv. to  $M$ .





(a)




(b)

[C. Adams, 1986]

Let  $L_1$  and  $L_2$  be links in  $S^3$  such that  $S^3 \setminus L_1$  and  $S^3 \setminus L_2$  are hyperbolic and  $L_1$  and  $L_2$  have projections as in Figure (a). Let  $L$  be the link with projection as in Figure (b). Then  $S^3 \setminus L$  is hyperbolic and


$$\text{vol}(S^3 \setminus L) = \text{vol}(S^3 \setminus L_1) + \text{vol}(S^3 \setminus L_2).$$

## Example 1




Tetrahedral manifold  $S^3 \setminus L_{8a20}$ .

## Example 2



Tetrahedral manifold = 3-fold covering of  $S^3 \setminus L8a20$ .

## Example 3



Almost tetrahedral manifold  $S^3 \setminus L10n84$ .



Thank you!