

О глобальных решениях полугамильтоновых систем квазилинейных уравнений

А.Е. Миронов

совместно с С.В. Агаповым (Новосибирск) и М. Бялым (Тель-Авив)

Декабрьские чтения в Томске
11-16 декабря 2018 г.

Quasi-linear PDEs

Quasi-linear systems of the form

$$A(U)U_x + B(U)U_y = 0,$$

$$U_t = A(U)U_x, \quad U = (u_1, \dots, u_n)^T$$

appears in such areas like

- gas-dynamics
- non-linear elasticity
- integrable geodesic flows on 2-torus


and many others.

Hopf equation

Consider the following equation $u_t + uu_x = 0$. The solution of the Cauchy problem $u|_{t=0} = g(x)$ is given by the implicit formula

$$u(x, t) = g(x - ut).$$

The solution becomes many valued.



- **Integrable geodesic flows**
- **Non-existence of non-trivial global solutions (smooth periodic) of quasi-linear systems:** *Hopf equations, model of non-linear elasticity, geodesic flows on 2-torus in elliptic domains*
- **Existence of non-trivial global solutions (smooth periodic) of quasi-linear systems:** *weakly non-linear systems, magnetic geodesic flows on 2-torus on one energy level*

Integrable geodesic flow on the 2-torus

Let

$$ds^2 = g_{ij}(x)dx^i dx^j, \quad i, j = 1, 2$$

be a Riemannian metric on \mathbb{T}^2 . The geodesic flow is called *integrable* if the Hamiltonian system

$$\dot{x}^i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial x^i}, \quad H = \frac{1}{2}g^{ij}p_i p_j$$

possesses an additional first integral $F : T^*\mathbb{T}^2 \rightarrow \mathbb{R}$ such that

$$\dot{F} = \{F, H\} = \sum_{j=1}^2 \left(\frac{\partial F}{\partial x^j} \frac{\partial H}{\partial p_j} - \frac{\partial F}{\partial p_j} \frac{\partial H}{\partial x^j} \right) = 0$$

and F is functionally independent with H almost everywhere.

Integrable geodesic flow on the 2-torus

Theorem (Bialy, M.)

If the Hamiltonian system has an integral F which is a homogeneous polynomial of degree n , then on the covering plane \mathbb{R}^2 there exist the global semi-geodesic coordinates (t, x) such that

$$ds^2 = g^2(t, x)dt^2 + dx^2, \quad H = \frac{1}{2}\left(\frac{p_1^2}{g^2} + p_2^2\right)$$

and F can be written in the form:

$$F_n = \sum_{k=0}^n \frac{a_k(t, x)}{g^{n-k}} p_1^{n-k} p_2^k.$$

Here the last two coefficients can be normalized by the following way:

$$a_{n-1} = g, \quad a_n = 1.$$

Integrable geodesic flow on the 2-torus

The condition $\{F, H\} = 0$ is equivalent to the quasi-linear PDEs

$$U_t + A(U)U_x = 0, \quad (1)$$

where $U^T = (a_0, \dots, a_{n-1})$, $a_{n-1} = g$,

$$A = \begin{pmatrix} 0 & 0 & \dots & 0 & 0 & a_1 \\ a_{n-1} & 0 & \dots & 0 & 0 & 2a_2 - na_0 \\ 0 & a_{n-1} & \dots & 0 & 0 & 3a_3 - (n-1)a_1 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{n-1} & 0 & (n-1)a_{n-1} - 3a_{n-3} \\ 0 & 0 & \dots & 0 & a_{n-1} & na_n - 2a_{n-2} \end{pmatrix}.$$

Semi-Hamiltonian systems

Theorem (Bialy, M.)

(1) is semi-Hamiltonian system. Namely, there is a regular change of variables

$$U \mapsto (G_1(U), \dots, G_n(U))$$

such that for some $F_1(U), \dots, F_n(U)$ the following conservation laws hold:

$$(G_i(U))_x + (F_i(U))_y = 0, \quad i = 1, \dots, n.$$

Moreover, in the hyperbolic domain, where eigenvalues $\lambda_1, \dots, \lambda_n$ of $A(U)$ are real and pairwise distinct, there exists a change of variables

$$U \mapsto (r_1(U), \dots, r_n(U))$$

such that the system can be written in Riemannian invariants:

$$(r_i)_x + \lambda_i(r)(r_i)_y = 0, \quad i = 1, \dots, n.$$

Systems of hydrodynamical type

$$\partial_{r_k} \left(\frac{\partial_{r_i} \lambda_j}{\lambda_i - \lambda_j} \right) = \partial_{r_i} \left(\frac{\partial_{r_k} \lambda_j}{\lambda_k - \lambda_j} \right),$$

$$\Gamma_{ij}^i = \partial_{r_j} \log \sqrt{g_{ii}} = \left(\frac{\partial_{r_j} \lambda_i}{\lambda_j - \lambda_i} \right),$$

$$ds^2 = g_{11}(r)(dr_1)^2 + \cdots + g_{nn}(dr_n)^2.$$

Generalized hodograph method

$$\frac{\partial_{r_j} w_i}{w_i - w_j} = \frac{\partial_{r_j} \lambda_i}{\lambda_j - \lambda_i},$$

$$w_i = \lambda_i t + x, \quad i = 1, \dots, n.$$

Geodesic flow on the 2-torus in elliptic region

Theorem (Bialy, M.)

Let $n = 4$, then in the elliptic regions the following alternative holds: either metric is flat or F_4 is reducible, that is it can be expressed as:

$$F_4 = k_1 F_2^2 + 2k_2 H F_2 + 4k_3 H^2$$

where F_2 is a polynomial of degree 2 which is an integral of the geodesic flow in the elliptic region and k_i are constants.

Geodesic flow on the 2-torus in elliptic region

The following theorem is crucial in proof of the previous one.

Theorem (Bialy, M.)

Assume that $\Omega_e = \mathbb{T}^2$ and assume that for all (t, x) the polynomial G_4 has 4 distinct roots, 2 – complex conjugate $s_{1,2} = \alpha \pm i\beta$ and 2 real $s_{3,4}$. Assume also that the imaginary part of Riemann invariants $r_{1,2}$ does not vanish. Then the real eigenvalues $\lambda_{3,4} = gs_{3,4}$ are necessarily genuinely non-linear and therefore the corresponding Riemann invariants are constants. In particular all a_i must be constant, and so the metric is flat.

Model of non-linear elasticity

Consider the following equation:

$$u_{tt} + (\sigma(u))_{xx} = 0, \quad u(t, x + 1) = u(t, x). \quad (1)$$

It can be viewed as a compatibility condition of the quasi-linear system of the form:

$$\begin{aligned} u_t &= -v_x, \\ v_t &= (\sigma(u))_x. \end{aligned} \quad (2)$$

Model of non-linear elasticity

Theorem (Bialy, M.)

If the function $\sigma(u)$ is either of quadratic-like or cubic-like type, then any C^2 -solution $(u(t, x), v(t, x))$ of the system (2) defined on the half-cylinder $[t_0, +\infty) \times \mathbb{S}^1$ so that

$$u(t, x + 1) = u(t, x), \quad v(t, x + 1) = v(t, x), \quad t \geq t_0,$$

which has initial values in the Hyperbolic region $U_h = \{u < \alpha\} \cup \{u > \beta\}$ must be constant.

Theorem (Bialy, M.)

If the function $\sigma(u)$ is either of quadratic-like or cubic-like type, then any C^2 -solution $(u(t, x), v(t, x))$ of the system (2) defined on the whole cylinder $\mathbb{R} \times \mathbb{S}^1$ so that

$$u(t, x + 1) = u(t, x), \quad v(t, x + 1) = v(t, x)$$

must be constant.

Model of non-linear elasticity

These theorems follow from the following facts. The system (2) can be written in the form:

$$\begin{pmatrix} u \\ v \end{pmatrix}_t + A(u, v) \begin{pmatrix} u \\ v \end{pmatrix}_x = 0, \quad A = \begin{pmatrix} 0 & 1 \\ -\sigma'(u) & 0 \end{pmatrix}.$$

In the hyperbolic region U_h the matrix A has two real distinct eigenvalues:

$$\lambda_1 = \sqrt{-\sigma'(u)}, \quad \lambda_2 = -\sqrt{-\sigma'(u)}.$$

Riemannian invariants have the form:

$$r_1 = v - \int_u^\alpha \sqrt{-\sigma'(u)} du, \quad r_2 = v + \int_u^\alpha \sqrt{-\sigma'(u)} du,$$
$$(r_i)_t + \lambda_i (r_i)_x = 0, \quad i = 1, 2.$$

Model of non-linear elasticity

The crucial fact here is that both eigenvalues are genuinely non-linear in U_h by the formulas:

$$(\lambda_1)_{r_1} = (\lambda_2)_{r_2} = \frac{\sigma''(u)}{4\sigma'(u)} \neq 0.$$

Along characteristics we get the following Riccati equations:

$$L_{v_1}(z_1) + kz_1^2 = 0, \quad L_{v_2}(z_2) + kz_2^2 = 0,$$

where

$$z_1 = (r_1)_x(-\sigma'(u))^{\frac{1}{4}}, \quad z_2 = (r_2)_x(-\sigma'(u))^{\frac{1}{4}}, \quad k = -\frac{\sigma''}{4(-\sigma'(u))^{\frac{5}{4}}}$$

and

$$L_{v_1} = \partial_t + \lambda_1 \partial_x, \quad L_{v_2} = \partial_t + \lambda_2 \partial_x$$

in what follows non-existence of periodic non-constant solutions.

Magnetic geodesic flow

Consider Hamiltonian system

$$\dot{x}^j = \{x^j, H\}_{mg}, \quad \dot{p}_j = \{p_j, H\}_{mg}, \quad j = 1, 2$$

on a 2-torus in magnetic field with $H = \frac{1}{2}g^{ij}p_i p_j$ and the Poisson bracket:

$$\{F, H\}_{mg} = \sum_{i=1}^2 \left(\frac{\partial F}{\partial x^i} \frac{\partial H}{\partial p_i} - \frac{\partial F}{\partial p_i} \frac{\partial H}{\partial x^i} \right) + \Omega(x^1, x^2) \left(\frac{\partial F}{\partial p_1} \frac{\partial H}{\partial p_2} - \frac{\partial F}{\partial p_2} \frac{\partial H}{\partial p_1} \right). \quad (6)$$

Example

Let $ds^2 = \Lambda(y)(dx^2 + dy^2)$ and the magnetic form $\omega = -u'(y)dx \wedge dy$. Then the magnetic geodesic flow is integrable and the first integral is linear in momenta:

$$F_1 = p_1 + u(y).$$

Main result

Theorem

There exist real analytic Riemannian metrics on the 2-torus which are arbitrary close to the Liouville metrics (and different from them) and a non-zero analytic magnetic fields such that magnetic geodesic flows on the energy level $\{H = \frac{1}{2}\}$ have polynomial in momenta first integral of degree two.

Magnetic geodesic flow

Choose the conformal coordinates (x, y) , such that $ds^2 = \Lambda(x, y)(dx^2 + dy^2)$, $H = \frac{p_1^2 + p_2^2}{2\Lambda}$. On the fixed energy level $H = \frac{1}{2}$ one can parameterize momenta by the following way:

$$p_1 = \sqrt{\Lambda} \cos \varphi, \quad p_2 = \sqrt{\Lambda} \sin \varphi.$$

Hamiltonian equations take the form

$$\dot{x} = \frac{\cos \varphi}{\sqrt{\Lambda}}, \quad \dot{y} = \frac{\sin \varphi}{\sqrt{\Lambda}}, \quad \dot{\varphi} = \frac{\Lambda_y}{2\Lambda\sqrt{\Lambda}} \cos \varphi - \frac{\Lambda_x}{2\Lambda\sqrt{\Lambda}} \sin \varphi - \frac{\Omega}{\Lambda}.$$

We shall search F in the form

$$F(x, y, \varphi) = \sum_{k=-N}^{k=N} a_k(x, y) e^{ik\varphi}. \quad (7)$$

Here $a_k = u_k + iv_k$, $a_{-k} = \bar{a}_k$. Condition $\dot{F} = \{F, H\}_{mg} = 0$ is equivalent to the following equation

$$F_x \cos \varphi + F_y \sin \varphi + F_\varphi \left(\frac{\Lambda_y}{2\Lambda} \cos \varphi - \frac{\Lambda_x}{2\Lambda} \sin \varphi - \frac{\Omega}{\sqrt{\Lambda}} \right) = 0. \quad (8)$$

Magnetic geodesic flow

Substituted (7) to (8), all the coefficients of $e^{ik\varphi}$ must be equal to zero. One obtains

$$\begin{aligned} \frac{\Lambda_y}{2\Lambda} \frac{i(k-1)a_{k-1} + i(k+1)a_{k+1}}{2} - \frac{\Lambda_x}{2\Lambda} \frac{i(k-1)a_{k-1} - i(k+1)a_{k+1}}{2i} + \\ + \frac{(a_{k-1})_x + (a_{k+1})_x}{2} + \frac{(a_{k-1})_y - (a_{k+1})_y}{2i} - \frac{ik\Omega a_k}{\sqrt{\Lambda}} = 0, \end{aligned} \quad (9)$$

where $k = 0, \dots, N+1$, $a_k = 0$ while $k > N$.

One can eliminate Ω from this system thus obtaining the quasilinear PDEs on a_j and Λ of a kind

$$A(U)U_x + B(U)U_y = 0, \quad (10)$$

where $U = (\Lambda, u_0, u_1, \dots, u_{N-1}, v_1, \dots, v_{N-1})^T$.

Semi-Hamiltonian system

Let $N = 2$. Then (10) takes the form

$$A(U)U_x + B(U)U_y = 0,$$

where

$$A = \begin{pmatrix} 0 & 0 & 1 & 0 \\ f & 0 & \Lambda & 0 \\ 2 & 1 & 0 & \frac{g}{2} \\ 0 & 0 & 0 & -\frac{f}{2} \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 & 1 & 0 \\ -g & 0 & 0 & -\Lambda \\ 0 & 0 & -\frac{g}{2} & 0 \\ 2 & -1 & \frac{f}{2} & 0 \end{pmatrix}, \quad (11)$$

$$U = (\Lambda, u_0, f, g)^T, \quad f = \frac{u_1}{\sqrt{\Lambda}}, \quad g = \frac{v_1}{\sqrt{\Lambda}}.$$

Magnetic field takes the form: $\Omega = \frac{1}{4}(g_x - f_y)$.

Crucial construction

One can check that

$$U_0(x, y) = \begin{pmatrix} \Lambda_1(x) + \Lambda_2(y) \\ 2\Lambda_2(y) - 2\Lambda_1(x) \\ 0 \\ 0 \end{pmatrix} \quad (12)$$

is the solution of the system (11), where $\Lambda_1(x)$ and $\Lambda_2(y)$ are periodic positive functions: $\Lambda_1(x+1) = \Lambda_1(x)$, $\Lambda_2(y+1) = \Lambda_2(y)$. This solution corresponds to the case of geodesic flow of the Liouville metric with zero magnetic field having the first integral of the second degree of the form

$$F_2 = \frac{\Lambda_2(y)p_1^2 - \Lambda_1(x)p_2^2}{\Lambda_1(x) + \Lambda_2(y)}.$$

Λ_1 and Λ_2 are assumed to be real analytic functions.

Introduce the following evolution equations:

$$U_t = A_1(U)U_x + B_1(U)U_y, \quad (13)$$

where

$$A_1 = \begin{pmatrix} g & 0 & 0 & \Lambda \\ -2g & g & 0 & -2\Lambda \\ 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \end{pmatrix}, \quad B_1 = \begin{pmatrix} f & 0 & \Lambda & 0 \\ 2f & f & 2\Lambda & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

This system defines the symmetry of the system (11) so that the flow of (13) transforms solutions to solutions as we shall prove below.

Next we apply the following consequence of Cauchy–Kowalevskaya theorem:

Lemma

The Cauchy problem for the system (13) with the initial data

$$U(x, y, t) |_{t=0} = U_0(x, y) \quad (14)$$

has a unique analytic periodic ($U(x + 1, y, t) = U(x, y + 1, t) = U(x, y, t)$) solution for t small enough.

Let us prove that $U(x, y, t)$ constructed in Lemma 1 is a solution of our system (9) for all small t . We denote by $\tilde{V}(x, y, t)$ the following real analytic vector function

$$\tilde{V} = A(U)U_x + B(U)U_y.$$

By our construction $\tilde{V}(x, y, 0) = 0$. We have to prove that $\tilde{V} \equiv 0$. Denote

$$\tilde{V} = \begin{pmatrix} V_1 \\ V_2 \\ V_3 \\ V_4 \end{pmatrix}, \quad V = \begin{pmatrix} V_2 \\ V_3 \\ V_4 \end{pmatrix}.$$

By direct calculations using (13) one can check that \tilde{V} satisfies the following system of equations:

$$\begin{pmatrix} V_1 \\ V_2 \\ V_3 \\ V_4 \end{pmatrix}_t = A_2 \begin{pmatrix} V_1 \\ V_2 \\ V_3 \\ V_4 \end{pmatrix}_x + B_2 \begin{pmatrix} V_1 \\ V_2 \\ V_3 \\ V_4 \end{pmatrix}_y + C_2 \begin{pmatrix} V_1 \\ V_2 \\ V_3 \\ V_4 \end{pmatrix} + D_2 \begin{pmatrix} V_1^2 \\ V_1 V_2 \\ V_1 V_3 \\ V_2 V_3 \end{pmatrix}. \quad (15)$$

Here

$$A_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ -g\Lambda & g & 0 & -2\Lambda \\ 0 & 0 & 0 & 0 \\ 2\Lambda & -2 & f & g \end{pmatrix}, \quad B_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ f\Lambda & f & 2\Lambda & 0 \\ 2\Lambda & 2 & f & g \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$C_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ c_1 & c_2 & c_3 & -2\Lambda_x \\ c_4 & 0 & f_y & -f_x \\ c_5 & 0 & -g_y & g_x \end{pmatrix}, \quad D_2 = \begin{pmatrix} 0 & 0 & 0 & 0 \\ -\frac{f\Lambda}{g} & -\frac{f}{g} & -\frac{4\Lambda}{g} & -\frac{4}{g} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

where

$$c_1 = \frac{2\Lambda f_x f + (f^2 - g^2 + 4\Lambda)\Lambda_x + 2\Lambda u_{0x}}{g}, \quad c_2 = \frac{2(gg_x + 2\Lambda_x + u_{0x})}{g},$$

$$c_3 = \frac{4\Lambda f_x + 2f\Lambda_x}{g}, \quad c_4 = 4\Lambda_y + \frac{1}{2}f(f_y - g_x), \quad c_5 = 4\Lambda_x - \frac{1}{2}g(f_y - g_x).$$

References

- [1]. Agapov S.V., Bialy M., Mironov A.E.: Integrable magnetic geodesic flows on 2-torus: new examples via quasi-linear system of PDEs. CMP, 2017. Vol. 351. N. 3. P. 993–1007.
- [2]. Bialy M.L., Mironov A.E.: New semi-hamiltonian hierarchy related to integrable magnetic flows on surfaces, Cent. Eur. J. Math., **10**:5, 1596–1604 (2012).
- [3]. Bialy M.L., Mironov A.E.: Rich quasi-linear system for integrable geodesic flow on 2-torus, Discrete and Continuous Dynamical Systems - Series A, **29**:1, 81–90 (2011).
- [4]. Bialy M.L., Mironov A.E.: Integrable geodesic flows on 2-torus: formal solutions and variational principle, J. Geom. and Phys., **87**:1, 39–47 (2015).
- [5]. Bialy M.L., Mironov A.E.: Cubic and quartic integrals for geodesic flow on 2-torus via a system of the hydrodynamic type, Nonlinearity, **24**:12, 3541–3554 (2011).

References

[6]. Bialy M.L., Mironov A.E.: From polynomial integrals of Hamiltonian flows to a model of non-linear elasticity, *Journal of Differential Equations*, **255**:10, 3434–3446 (2013).